

1 Title page

2
3 Effective Brief, Low-impact, High-intensity Osteogenic Loading in Postmenopausal
4 Osteoporosis

5
6 Disclosure 1: Professor George Chrousos is an Editorial Board Member for The
7 Journal of Clinical Endocrinology & Metabolism and played no role in the Journal's
8 evaluation of the manuscript.

9
10 Disclosure 2: The study was performed with funding provided to the University
11 Research Institute of Maternal and Child Health and Precision Medicine, National
12 and Kapodistrian University of Athens, Athens, Greece, by Osteofit Hellas, Inc.

13
14 Short running head:

15
16 Osteostrong® use in Osteoporosis

17
18
19
20
21 Nektaria Papadopoulou–Marketou^{1,2}, Anna Papageorgiou², Nikolaos Marketos³,
22 Panagiotis Tsiamyrtzis^{4,5}, Georgios Vavetsis², George P. Chrousos²

23
24 ¹Neuroendocrine Tumor Unit, ENETS Centre of Excellence, 1st Department of
25 Propaedeutic and Internal Medicine, Laiko General Hospital, National and
26 Kapodistrian University of Athens, 11527 Athens, Greece

27
28 ²University Research Institute of Maternal and Child Health and Precision
29 Medicine, National and Kapodistrian University of Athens, Athens, Greece

³Department of Physiology, Medical School, National and Kapodistrian University of Athens, Athens, Greece

⁴Dept. of Mechanical Engineering, Politecnico di Milano, Italy

⁵Dept. of Statistics, Athens University of Economics and Business, Greece.

Corresponding author: Nektaria Papadopoulou–Marketou, MD, PhD,
Neuroendocrine Tumor Unit, ENETS Centre of Excellence, 1st Department of
Propaedeutic and Internal Medicine, Laiko General Hospital, National and
Kapodistrian University of Athens, 11527 Athens, Greece

E-mail: papadopoulounektaria@gmail.com

ORCID ID: <https://orcid.org/0000-0002-6719-7923>

Keywords: Osteostrom, Osteoporosis, Bone metabolism, Exercise in
osteoporosis, Osteogenic loading, antiosteoporotic treatment

Abstract

Background: Osteoporosis is characterized by reduced bone mineral density (BMD) and disrupted microarchitecture estimated by trabecular bone score (TBS), resulting in increased bone fracture risk. "Osteostrom[®]" is a bone-strengthening system implementing 4 devices and incorporating brief (10-minute), weekly, low-

1 impact, and high-intensity osteogenic loading exercises. We evaluated the efficacy
2 of the Osteostrong® intervention in postmenopausal osteoporotic women.

3 Methods and Subjects:

4 147 postmenopausal osteoporotic women were separated into two groups: Group
5 A comprised 74 women receiving Osteostrong® intervention (mean age 58.8 years,
6 56.6-61 years 95% CI), and was subdivided into G1 receiving no antiresorptive
7 medication, and G2 on such medication. Group B comprised 73 women that
8 received no Osteostrong® intervention (mean age 61.8 years, 59.4-64.1 95% CI)
9 and was subdivided into G3 on no antiresorptive therapy, and G4 on such
10 treatment. All participants underwent a physical examination and had an
11 assessment for secondary osteoporosis. Dual-energy X-ray absorptiometry (DXA)
12 examinations (Horizon W [S/N 300472M]) were performed at the time of trial
13 inclusion and 12 months later.

14 Results: Statistically significant increases were observed in the following
15 parameters: i) BMD of the lumbar spine (L1-L4) in G1($p=0.0039$), G2($p<0.001$),
16 and G4($p=0.0059$): ii) TBS in G2($p=0.0078$): iii) BMD of the right femoral neck in
17 G1($p=0.0382$) and G4($p=0.032$): iv) BMD of the left femoral neck G2($p=0.0089$)
18 and in G4($p=0.0498$) and total femur in G2($p=0.0162$).

19 Conclusions: Osteostrong® improved BMD of the lumbar spine in women with
20 osteoporosis both off and on antiresorptive treatment. Furthermore, Osteostrong®
21 enhanced the effect of antiresorptive therapy on BMD and TBS of the spine, hip
22 and femoral neck.

1

2

3 **Introduction**

4 At the dawn of the 21st century, health and wellness-directed measures in Western
5 civilization seem to have successfully addressed many problems of everyday life.

6 Nevertheless, an increased fracture risk state, known as osteoporosis, which is
7 characterized by decreased bone mineral density (BMD) and disrupted
8 microarchitecture due to various factors, remains a global health burden. The
9 endpoint of osteoporosis is the low-energy fracture, and pain, immobilization, and
10 disability, in general, remain some of the main reasons for drug development and
11 research on alternative methods to delay or postpone their incidence. Expected
12 fragility fractures per year rise to 4.5 million in the European Union and 1.5 million
13 in the USA, at a 12- to 18-billion-dollar expense for the healthcare systems with a
14 previous report projecting even higher costs in the years to come (1, 2). Almost $\frac{3}{4}$
15 of these fractures affect postmenopausal women. This makes the prevention of
16 osteoporotic fractures a global priority for healthcare systems.

17 Apart from dual-energy x-ray absorptiometry (DXA), which is currently considered
18 the gold standard for evaluating BMD, measurement of the trabecular bone score
19 (TBS) that consists of a grey-level textural measurement, which is typically
20 obtained from conventional lumbar spine DXA- BMD images, serves as a validated
21 index of bone microarchitecture that is correlated with the mechanical properties
22 of bone (3). TBS has been added to the already established methods of fracture
23 risk assessment tool (FRAX) and BMD for predicting fracture risk (3).

Despite the plethora of drugs used to delay and/or treat osteoporosis, very few non-pharmaceutical methods have been tested and even fewer have succeeded. Until recently, the treatment of osteoporosis in the setting of bone fractures, especially those of the lumbar spine level, consisted of pharmacotherapy, surgery, and the avoidance of everyday exercise, with controversial results. The literature describes non-pharmaceutical methods to reduce fracture risk due to osteoporosis as traditional, complementary, and integrative medicine (4). Patients often use these methods without informing their physicians, making it even more difficult to evaluate the possible protective effects of these practices against osteoporosis (4). Among these proposed methods, supplementation of calcium and vitamin D along with exercise remains the most often recommended modality for osteoporosis prevention according to worldwide clinical practice guidelines, although specific parameters of exercise are not defined (4-6).

There is substantial patient interest in non-pharmacologic approaches to treating or preventing osteoporosis and Osteostrom® is one such intervention, whose efficiency, however, has not been evaluated in clinical trial yet. Interestingly, suggestions for exercise to strengthen the musculoskeletal system and to increase balance have been provided in recent treatment guidelines for osteoporosis and fracture prevention, without specific instructions for the duration, extension, type of exercise, and other important parameters (7, 8).

Osteogenic loading is based on Wolff's law suggesting that a healthy animal's bone will adjust to the forces applied to it (9). The internal structure of trabeculae experiences adaptive alterations, which are subsequently followed by secondary adjustments to the external cortical part of the bone. This process could result in

1 the thickening of the bone. Conversely, when the load on a bone is reduced, the
2 bone experiences a decrease in density and strength due to the absence of the
3 necessary stimulus required for continuous remodeling (10-12). Osteostrong®, a
4 brief, low-impact, and high-intensity osteogenic loading training modality with
5 once-weekly, 10-minute sessions, using patented devices, known as “Spectrum”,
6 is described below. Although several people have used the Osteostrong®
7 intervention with reportedly beneficial effects on bone density, the method has
8 been less studied for its efficacy in improving bone density and quality in subjects
9 with osteopenia or osteoporosis.

10 **Methods and Study Population**

11 **Osteostrong® Method**

12 Osteostrong® consists of a series of brief, 10-min total, weekly exercises of
13 osteogenic loading tailored specifically for every person to ameliorate
14 osteoporosis. This exercise is facilitated using four different apparatuses and is
15 characterized by low impact but high intensity, making it more attractive for patients
16 to adhere to and to avoid possible risks of lumbar or other injury, especially for the
17 elderly and individuals with more severe osteoporosis. This equipment applies a
18 certain dosage of force specific to the participants’ multiples of bodyweight (MOB),
19 allowing for an increased likelihood of osteogenesis. Spectrum devices emulate
20 these forces in a safe and controlled environment. Designed to enable users to
21 assume optimal leverage positions, each device allows high-force application in
22 static postures, aiming to achieve bone-strengthening impact without actual high-
23 impact activity, taking advantage of the individual’s physiological strength limit
24 preventing injury. The four Spectrum devices are based on Growth Trigger (GT),

1 indicating each device's objective to facilitate bone adaptation (Upper GT, Lower
2 GT, Postural GT, and Core GT), each focusing on distinct regions of the
3 musculoskeletal system. The Upper GT targets the kinetic chain from hands to
4 clavicle; the Lower GT from feet to hip; the Postural GT from feet to neck; finally,
5 the Core GT focuses on the rib cage. Moreover, the Osteostrong® intervention
6 program includes falls prevention and balance training, with a focus on posture,
7 stability, mobility, and coordination.

8 **Study Population**

9 A total of 147 postmenopausal women with osteoporosis of the lumbar spine and/or
10 femoral neck or total hip, who were followed at and recruited from the Unit on
11 Clinical and Translational Research in Endocrinology, National and Kapodistrian
12 University of Athens, Greece, were enrolled. All participants were Caucasian adults
13 of Greek origin. A consent form was obtained from all the participants. Inclusion
14 criteria were female gender, menopause that presented either early or at the
15 expected age, osteoporosis measured by DXA in either lumbar spine and/or hip.
16 Exclusion criteria were secondary osteoporosis and recent bone fracture. Patients
17 were assigned to the various treatment groups according to their status and
18 preference, after they had been informed about the Osteostrong® intervention.
19 Patients were divided into two groups. Group A included 74 women treated with
20 Osteostrong® (mean age: 58.8y, 95%CI 56.6-61); Group A was subdivided into G1,
21 which included women who had no parallel antiresorptive treatment, and G2, which
22 included women who were treated in parallel with either oral bisphosphonates or
23 denosumab. Group B included 73 women who had no Osteostrong® intervention
24 (mean age 61.8y, 95%CI 59.4-64.1). Group B was subdivided into G3, which

1 included women who did not receive antiresorptive treatment, and G4, which
2 included women who were treated with such medication. Data regarding BMI and
3 age are shown in Table 1. None of the patients from either groups G1 or G3 had
4 previous antiresorptive treatment. Regarding groups G2 & G4, for detailed
5 information, see the supplementary Table S1 (13). In groups G2 and G4,
6 Denosumab, alendronate and risendronate were the antiresorptive drugs used. All
7 patients received 1200 mg calcium and Vitamin D supplementation as required.

8 All participants underwent a complete physical examination and an assessment
9 for exclusion of secondary osteoporosis. Details from medical history, such as
10 uptake of glucocorticoids or previous fragility bone fracture were retrieved, and
11 laboratory tests to exclude primary hyperparathyroidism, connective tissue
12 disorders, hypercortisolism, Vitamin D insufficiency or other conditions, such as
13 sarcoidosis or blood malignancy, were conducted. DXA examination [Horizon W
14 (S/N 130472M)] twice, at the time of inclusion in the trial and 12 months later, was
15 performed. Bone markers [C-terminal telopeptide of type I collagen (CTX-I) and N-
16 terminal propeptide of type I procollagen (PINP)] were examined after the
17 intervention in groups G1 and G2. Statistical analysis was performed using the
18 freeware R (4.2.2) and examined during the study period for significant mean
19 differences in the recorded response variables.

20 **Results**

21 The main adverse event that was feared by the participants and closely monitored
22 by the Osteostrom® facility personnel was a new bone fracture at any site and/or
23 a muscle or tendon trauma during the procedure. None of these events occurred
24 to any participant. In total, 13 patients experienced joint pain, and 3 patients

1 dropped out from the study because of it. BMD measurements at baseline and 12
 2 months later are shown in Table 2. Paired Student t-test of all the parameters
 3 before and after the Osteostrom® intervention showed the following: a statistically
 4 significant increase in T-score (L1-L4) and BMD of the lumbar spine (L1-L4) in
 5 groups G1 (n=50, % of positive differences (End-Start) = 72 %, Mean difference
 6 0.245, p<0.001) and (n=48, % of positive differences (End-Start): 56.2%,
 7 p<0.0039) respectively, G2 (n=21, % of positive differences (End-Start) = 85.7%,
 8 p<0.001) and (n=21, % of positive differences (End-Start) = 76.2%, p<0.001)
 9 respectively, and G4 (n=25, % of positive differences (End-Start) = 72%, p=0.0024)
 10 and (n=26, % of positive differences (End-Start) = 65.4%, p=0.0059), respectively.
 11 A statistically significant increase in T-score (TBS), as well as TBS, was found in
 12 the G2 group (n=18, % of positive differences (End-Start) = 66.7%, p=0.0025) and
 13 (n=20, % of positive differences (End-Start) = 60%, p=0.0078), respectively. A
 14 statistically significant increase in T-score and BMD of the right femoral neck BMD
 15 was shown in G1 group (n=44, % of positive differences (End-Start) = 47.7%,
 16 p=0.0496) and (n=44, % of positive differences (End-Start) = 47.7%, p=0.0382),
 17 respectively. A statistically significant increase in T-score and BMD of the Left
 18 femoral neck was shown in G2 group (n=22, % of positive differences (End-Start)
 19 = 54.5%, p=0.0152) and (n=22, % of positive differences (End-Start) = 63.6%,
 20 p=0.0089) respectively. A statistically significant increase in T-score and BMD of
 21 the Left femur total was shown in G2 group (n=22, % of positive differences (End-
 22 Start) = 54.5, p=0.0162) and (n=21, % of positive differences (End-Start) = 66.7%,
 23 p=0.0304), respectively, and G4 but only regarding BMD (n=27, % of positive
 24 differences (End-Start) = 70.4%, p=0.0498). All data are shown in Figure 1 and
 25 Table 2. Regression analysis regarding age and BMI was assessed, but no

1 statistically significant results were observed, Table S2 (13). For G2 and G4
2 patients receiving Denosumab, a regression analysis regarding TBS changes was
3 performed, but although a trend for increased TBS was observed, it was not
4 statistically significant (Table S3, Figure S1)(13) . For G3 patients who received
5 neither Osteostrom[®] intervention nor antiresorptive treatment a slight deterioration
6 was seen in all parameters but none of them was statistically significant. No
7 statistically significant correlations were found in either P1NP or CTX-1 levels and
8 BMD changes, in any of the groups.

9 **Discussion**

10 This is the first study that clearly demonstrates benefit from Osteostrom[®]
11 intervention in postmenopausal women. A significant improvement in BMD and T-
12 score in the lumbar spine in subjects who had an Osteostrom[®] intervention
13 regardless of antiresorptive medication was found. Additionally, the TBS of the
14 spine was significantly improved in the group that received the combination of
15 Osteostrom[®] intervention and anti-osteoporotic medication. Moreover, BMD and
16 T-score of the femoral neck and hip in both left and right side improved significantly
17 in subjects who had an Osteostrom[®] intervention and furthermore in those who
18 also had anti-osteoporotic treatment. As expected, patients without any
19 pharmaceutical or training intervention showed a slight, non-significant decline in
20 all mentioned bone parameters.

21 Osteoporosis reflects the risk of bone fracture, often resulting in pain, surgery, and
22 complications owing to prolonged hospitalization and workforce loss. Osteoporosis
23 is a global socioeconomic burden that affects millions of people and is associated
24 with a low quality of life. Despite the administration of numerous treatment

1 modalities over the years, no nonpharmacological approach has demonstrated
2 satisfactory results.

3 Literature has extensively discussed various exercise modalities and their effects
4 on bone mass and, eventually, osteoporosis. The notion of a “mechanostat” on
5 bone was introduced by Wolff and then described in detail by Frost (9-12). The
6 idea of a bone sensor that responds positively to external forces through muscle
7 training, resulting in improved thickness and microarchitecture has been
8 compelling but never shown in a clinical trial. This is the first time that the
9 administration of training in a calculated and individually tailored manner has
10 shown a positive effect on the density of the bone marrow of the spine. The lack of
11 contraindications to this treatment modality, as it does not interfere with any other
12 medication, makes it more attractive irrespective of age, socioeconomic status,
13 and comorbidities, resulting in strong adherence to an exercise plan.

14 Earlier studies have shown a positive effect of low-load, high-repetition resistance
15 exercise on the lumbar spine BMD of otherwise healthy postmenopausal women
16 compared to controls who did not exercise (14). Although the beneficial effect of
17 progressive resistance training on the lumbar spine was not indicated in any of the
18 literature reviews (15), perhaps because of the large variation in the studies
19 included, the approach of low-intensity, high-repetitive exercise tends to be
20 abandoned as more recent evidence emerges. Alternative methods of training,
21 such as Tai Chi Chuan, have also shown beneficial effects on lumbar spine BMD
22 (16), but availability is certainly an issue. In men with osteopenia, High-intensity
23 resistance, and impact training (HiRIT) has shown a beneficial effect on bone
24 geometry and strength in the femoral neck compared to isometric training based

on machines (17). Similarly, stable values for lumbar spine BMD after 12 months of high-intensity exercise were shown in men with osteosarcopenia according to the Franconian Osteopenia and Sarcopenia Trial (FrOST) (18).

Previous studies have shown that osteogenic loading positively affects spinal osteoporosis. HiRIT has been assessed previously in postmenopausal women with encouraging results (17-23). Additionally, the high-intensity exercise implemented not only did not cause extra-vertebral fractures but also improved thoracic kyphosis in this group of patients (17, 20, 22). Moreover, HiRIT was associated with greater improvement in BMD of the lumbar spine of osteoporotic postmenopausal women compared with medium-intensity resistance and impact training (MiRIT), thus diminishing the fear of new fractures in this sensitive population, despite the small number of clinical trials evaluating HiRIT (23-26).

According to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (27), literature reviews based on (a) controlled trials, (b) isolated DRT with at least one exercise and one control group, (c) intervention durations ≥ 6 months, (d) BMD assessments at the lumbar spine or proximal femur, and (e) cohorts of postmenopausal women, revealed that once-weekly exercise benefitted postmenopausal women with osteoporosis in comparison to more frequent exercise concerning lumbar spine BMD, and was the sole parameter that could be suggested regarding the modus of exercise (25, 26, 28-30). Similar positive results for lumbar spine BMD after high-intensity exercise were reported in another review further supporting this argument (31). Moreover, positive effects of high-intensity exercise on lumbar spine BMD in postmenopausal women with osteopenia or osteoporosis were also shown in the ACTLIFE-RCT trial

(32). Nevertheless, it is difficult to address all parameters needed for optimal outcomes through exercise. Even when classified according to the type of exercise (weight-bearing, dynamic resistance, and mixed interventions), no clear differences were observed, as all were beneficial for the lumbar spine BMD of postmenopausal women (29). A recent review and meta-analysis isolated 14 of 780 studies concerning the mode (resistance only vs. combined resistance and weight-bearing exercises), frequency, volume, load, and program length (33). It was shown that increases in BMD were favored by combined resistance and weight-bearing exercises, lower volumes, and higher loads, although lumbar spine BMD did not benefit from current resistance exercise programs, leaving room for improvement in this domain (33). To address the difficult aspects of the optimal resistance training modality, a new trial is currently underway (34).

The beneficial effect of progressive resistance training with specifically tailored, multicomponent exercise programs, along with other health adjustments for balance and nutrition, have been recommended for the management of patients with osteoporosis. Implementation of these encouraging results fueled the recent UK guidelines. Resistance and impact training is recommended for patients with osteoporosis; however, it is clearly stated that the benefit of physical activity often outweighs the risk of harm (28). Nevertheless, a more careful approach regarding the intensity of exercise was adopted because of the lack of evidence of the beneficial effect of high- vs. moderate-intensity training.

Exercise recommendation for osteoporosis is often a sensitive matter, especially in the lumbar spine, as the incorrect application of force by the individual may result in new fractures, causing more pain and disability. Osteostrom[®], which consists of

1 a series of brief, 10-min, weekly exercises of osteogenic loading tailored
2 specifically for every person to ameliorate osteoporosis, showed opposite results.
3 It was well-tolerated by patients of different ages and socioeconomic levels who
4 followed this program for an entire year. Additionally, despite the intensity of the
5 exercise, no adverse events such as new fractures, muscle trauma, or other
6 injuries were reported, thus enhancing patient commitment to the exercise
7 program. This exercise is facilitated using four different apparatuses and is
8 characterized by low impact but high intensity, making it more attractive for patients
9 to adhere to and avoid possible risks of lumbar injury, especially for the elderly and
10 individuals with more severe osteoporosis. A previous pilot study on just one
11 subject, who was an astronaut, of weekly 15-min exercise for 6 months showed an
12 increase in body strength but no other osteoporosis parameters, perhaps due to
13 the short period of assessment (35, 36).

14 Similar results were reported from another group, as HiRIT showed a beneficial
15 effect in total hip, femoral neck volumetric and geometric section modules in
16 comparison to a low intensity Pilates based exercise (LiPBE). This effect was
17 further augmented in patients that received antiresorptive treatment in addition to
18 HiRIT (24).

19 The beneficial results of exercise in diminishing fracture risk, the main endpoint of
20 all anti-osteoporotic interventions, was observed in a retrospective, observational
21 study of men (37). This study demonstrated that vigorous but not moderate
22 exercise resulted in a decreased hazard ratio for fracture risk in life-long athletes,
23 and this effect was progressively more evident with passing years. These results

1 encourage the future application of Osteostrom® principles in other groups of
2 individuals vulnerable for osteopenia or osteoporosis.

3 Our study had some limitations. Not all patients who were asked to take part in the
4 study accepted to participate with various explanations (difficult to adhere to the
5 weekly appointments due to long distance, work schedule, lack of time, other
6 responsibilities, fear of trauma). Also, inadvertently, some serum parameters, such
7 as CTX and NTX concentrations, were not obtained. Furthermore, some patients
8 failed to follow-through and remain in the study during the entire year of treatment.

9 On the other hand, the study had certain advantages. The patients were recruited
10 from both the public and private medical sector, thus reflecting a socioeconomically
11 balanced group. Measurements of BMD and TBS were performed in the same
12 DXA machine in all patients, decreasing the variability and increasing the validity
13 of the results. Furthermore, the patients were referred from the entire Athens
14 metropolitan area to the same principal investigators, who screened and evaluated
15 all the participants in the study, using a similar approach.

16 In conclusion, the study demonstrated that the Osteostrom® intervention is a safe,
17 brief method with a significant positive impact on the BMD and TBS of the lumbar
18 spine, hip and femoral neck, in women with postmenopausal osteoporosis. The
19 study suggests that Osteogenic loading has an additive/synergistic effect with anti-
20 osteoporotic medication, augmenting the latter's efficacy and resulting in improved
21 bone strength and quality, and, hopefully, a reduced risk of bone fractures.

22

23

1 **Acknowledgements**

2 The study was performed with funding provided to the University Research
3 Institute of Maternal and Child Health and Precision Medicine, National and
4 Kapodistrian University of Athens, Athens, Greece, by Osteofit Hellas, Inc.

5 **Data Availability Statement**

6 The authors confirm that the data supporting the findings of this study are available
7 within the article and supplementary materials (13) and can be available upon
8 request.

9

10

11 **References**

- 12 1. Li N, Cornelissen D, Silverman S, Pinto D, Si L, Kremer I, et al. An Updated
13 Systematic Review of Cost-Effectiveness Analyses of Drugs for Osteoporosis.
14 Pharmacoeconomics. 2021;39(2):181-209.
- 15 2. Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A.
16 Incidence, and economic burden of osteoporosis-related fractures in the United
17 States, 2005-2025. J Bone Miner Res. 2007;22(3):465-75.
- 18 3. Shevroja E, Lamy O, Kohlmeier L, Koromani F, Rivadeneira F, Hans D. Use
19 of Trabecular Bone Score (TBS) as a Complementary Approach to Dual-energy X-
20 ray Absorptiometry (DXA) for Fracture Risk Assessment in Clinical Practice. J Clin
21 Densitom. 2017 Jul-Sep;20(3):334-345

- 1 4. Kocijan R, Medibach A, Lechner L, Haschka J, Kocijan A, Kraus DA, et al.
2 Use of Complementary and Alternative Medicine in Patients with Rare Bone
3 Diseases and Osteoporosis. *Nutrients*. 2024;16(6).
- 4 5. Coronado-Zarco R, Olascoaga-Gomez de Leon A, Garcia-Lara A,
5 Quinzanos-Fresnedo J, Nava-Bringas TI, Macias-Hernandez SI.
6 Nonpharmacological interventions for osteoporosis treatment: Systematic review
7 of clinical practice guidelines. *Osteoporos Sarcopenia*. 2019;5(3):69-77.
- 8 6. Choi YJ, Oh HJ, Kim DJ, Lee Y, Chung YS. The prevalence of osteoporosis
9 in Korean adults aged 50 years or older and the higher diagnosis rates in women
10 who were beneficiaries of a national screening program: the Korea National Health
11 and Nutrition Examination Survey 2008-2009. *J Bone Miner Res*. 2012;27(9):1879-
12 86.
- 13 7. Camacho PM, Petak SM, Binkley N, Diab DL, Eldeiry LS, Farooki A, et al.
14 American Association of Clinical Endocrinologists/American College of
15 Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of
16 Postmenopausal Osteoporosis-2020 Update. *Endocr Pract*. 2020;26(Suppl 1):1-
17 46.
- 18 8. Watts NB, Camacho PM, Lewiecki EM, Petak SM, Force AAOGT.
19 American Association of Clinical Endocrinologists/American College of
20 Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of
21 Postmenopausal Osteoporosis-2020 Update. *Endocr Pract*. 2021;27(4):379-80.
- 22 9. Ruff C, Holt B, Trinkaus E. Who's afraid of the big bad Wolff?: "Wolff's law"
23 and bone functional adaptation. *Am J Phys Anthropol*. 2006;129(4):484-98.

- 1 10. Wolf JH. [Julius Wolff and his "law of bone remodeling"]. *Orthopade.*
2 1995;24(5):378-86.
- 3 11. Frost HM. Bone's mechanostat: a 2003 update. *Anat Rec A Discov Mol Cell*
4 *Evol Biol.* 2003;275(2):1081-101.
- 5 12. Tyrovolas JB. The "Mechanostat Theory" of Frost and the
6 OPG/RANKL/RANK System. *J Cell Biochem.* 2015;116(12):2724-9.
- 7 13. Papadopoulou-Marketou N, Papageorgiou A, Marketos N, Tsiamyrtzis P,
8 Vavetsis G, Chrousos G. Effective Brief, Low-impact, High-intensity Osteogenic
9 Loading in Postmenopausal Osteoporosis (Supplementary session)
- 10 14. Nicholson VP, McKean MR, Slater GJ, Kerr A, Burkett BJ. Low-Load Very
11 High-Repetition Resistance Training Attenuates Bone Loss at the Lumbar Spine in
12 Active Post-menopausal Women. *Calcif Tissue Int.* 2015;96(6):490-9.
- 13 15. El-Kotob R, Ponzano M, Chaput JP, Janssen I, Kho ME, Poitras VJ, et al.
14 Resistance training and health in adults: an overview of systematic reviews. *Appl*
15 *Physiol Nutr Metab.* 2020;45(10 (Suppl. 2)):S165-S79.
- 16 16. Wu HY, Wang YR, Wen GW, Tang ZY, Yu YQ, Zhang JR, et al. Tai Chi on
17 bone mineral density of postmenopausal osteoporosis: A protocol for systematic
18 review and meta-analysis. *Medicine (Baltimore).* 2020;99(36):e21928.
- 19 17. Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. A
20 Comparison of Bone-Targeted Exercise Strategies to Reduce Fracture Risk in
21 Middle-Aged and Older Men with Osteopenia and Osteoporosis: LIFTMOR-M
22 Semi-Randomized Controlled Trial. *J Bone Miner Res.* 2020;35(8):1404-14.
- 23 1.8 Kemmler W, Kohl M, Frohlich M, Jakob F, Engelke K, von Stengel S, et al.
24 Effects of High-Intensity Resistance Training on Osteopenia and Sarcopenia

1 Parameters in Older Men with Osteosarcopenia-One-Year Results of the
2 Randomized Controlled Franconian Osteopenia and Sarcopenia Trial (FrOST). J
3 Bone Miner Res. 2020;35(9):1634-44.

4 19. Watson SL, Weeks BK, Weis LJ, Harding AT, Horan SA, Beck BR. High-
5 Intensity Resistance and Impact Training Improves Bone Mineral Density and
6 Physical Function in Postmenopausal Women With Osteopenia and Osteoporosis:
7 The LIFTMOR Randomized Controlled Trial. J Bone Miner Res. 2018;33(2):211-
8 20.

9 20. Watson SL, Weeks BK, Weis LJ, Harding AT, Horan SA, Beck BR. High-
10 intensity exercise did not cause vertebral fractures and improves thoracic kyphosis
11 in postmenopausal women with low to very low bone mass: the LIFTMOR trial.
12 Osteoporos Int. 2019;30(5):957-64.

13 21. Bonnet S, Paulin R, Sutendra G, Dromparis P, Roy M, Watson KO, et al.
14 Dehydroepiandrosterone reverses systemic vascular remodeling through the
15 inhibition of the Akt/GSK3- β /NFAT axis. Circulation. 2009;120(13):1231-40.

16 22. Harding AT, Weeks BK, Lambert C, Watson SL, Weis LJ, Beck BR. Effects
17 of supervised high-intensity resistance and impact training or machine-based
18 isometric training on regional bone geometry and strength in middle-aged and
19 older men with low bone mass: The LIFTMOR-M semi-randomised controlled trial.
20 Bone. 2020;136:115362.

21 23. Kistler-Fischbacher M, Yong JS, Weeks BK, Beck BR. A Comparison of
22 Bone-Targeted Exercise With and Without Antiresorptive Bone Medication to
23 Reduce Indices of Fracture Risk in Postmenopausal Women With Low Bone Mass:

- 1 The MEDEX-OP Randomized Controlled Trial. *J Bone Miner Res.*
2 2021;36(9):1680-93.
- 3 24. Kistler-Fischbacher M, Yong JS, Weeks BK, Beck BR. High-Intensity
4 Exercise and Geometric Indices of Hip Bone Strength in Postmenopausal Women
5 on or off Bone Medication: The MEDEX-OP Randomised Controlled Trial. *Calcif*
6 *Tissue Int.* 2022;111(3):256-66.
- 7 25. Kitagawa T, Hiraya K, Denda T, Yamamoto S. A comparison of different
8 exercise intensities for improving bone mineral density in postmenopausal women
9 with osteoporosis: A systematic review and meta-analysis. *Bone Rep.*
10 2022;17:101631.
- 11 26. Kitsuda Y, Wada T, Noma H, Osaki M, Hagino H. Impact of high-load
12 resistance training on bone mineral density in osteoporosis and osteopenia: a
13 meta-analysis. *J Bone Miner Metab.* 2021;39(5):787-803.
- 14 27. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD,
15 et al. The PRISMA 2020 statement: An updated guideline for reporting systematic
16 reviews. *J Clin Epidemiol.* 2021;134:178-89.
- 17 28. Brooke-Wavell K, Skelton DA, Barker KL, Clark EM, De Biase S, Arnold S,
18 et al. Strong, steady, and straight: UK consensus statement on physical activity
19 and exercise for osteoporosis. *Br J Sports Med.* 2022;56(15):837-46.
- 20 29. Shojaa M, von Stengel S, Kohl M, Schoene D, Kemmler W. Effects of
21 dynamic resistance exercise on bone mineral density in postmenopausal women:
22 a systematic review and meta-analysis with special emphasis on exercise
23 parameters. *Osteoporos Int.* 2020;31(8):1427-44.

- 1 30. Shojaa M, Von Stengel S, Schoene D, Kohl M, Barone G, Bragonzoni L, et
2 al. Effect of Exercise Training on Bone Mineral Density in Post-menopausal
3 Women: A Systematic Review and Meta-Analysis of Intervention Studies. *Front*
4 *Physiol.* 2020;11:652.
- 5
6 31. Manaye S, Cheran K, Murthy C, Bornemann EA, Kamma HK, Alabbas M,
7 et al. The Role of High-intensity and High-impact Exercises in Improving Bone
8 Health in Postmenopausal Women: A Systematic Review. *Cureus.*
9 2023;15(2):e34644.
- 10 32. Hettchen M, von Stengel S, Kohl M, Murphy MH, Shojaa M, Ghasemikaram
11 M, et al. Changes in Menopausal Risk Factors in Early Postmenopausal
12 Osteopenic Women After 13 Months of High-Intensity Exercise: The Randomized
13 Controlled ACTLIFE-RCT. *Clin Interv Aging.* 2021;16:83-96.
- 14 33. O'Bryan SJ, Giuliano C, Woessner MN, Vogrin S, Smith C, Duque G, et al.
15 Progressive Resistance Training for Concomitant Increases in Muscle Strength
16 and Bone Mineral Density in Older Adults: A Systematic Review and Meta-
17 Analysis. *Sports Med.* 2022;52(8):1939-60.
- 18 34. Giangregorio LM, Bleakney RR, Brien S, Butcher SJ, Chan BCF, Chilibeck
19 PD, et al. Finding the Optimal Resistance Training Intensity for Your Bones:
20 Protocol for a Randomized Controlled Trial. *Phys Ther.* 2023;103(10).
- 21 35. JafariNasabian P. How exercise and dietary intervention affect the outcome
22 of osteosarcopenic obesity syndrome? *Journal of Functional Morphology and*
23 *Kinesiology*: MDPI Multidisciplinary Digital Publishing Institute; 2018. p. 31-.

1 36. Tsung A, Jupiter D, Jaquish J, Sibonga J. Weekly Bone Loading Exercise
2 Effects on Healthy Subjects Strength, Bone Density, and Bone Biomarkers. *Aerosp*
3 *Med Hum Perform.* 2021;92(3):201-6.

4 37. Korhonen MT, Kujala UM, Kettunen J, Korhonen OV, Kaprio J, Sarna S, et
5 al. Longitudinal Associations of High-Volume and Vigorous-Intensity Exercise With
6 Hip Fracture Risk in Men. *J Bone Miner Res.* 2022;37(8):1562-70.

7

8 **Table 1:** BMI and age in the different subgroups.

9

	G1		G2		G3		G4	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
age	57.71	7.86	61.05	12.07	58.78	9.83	64.38	9.36
BMI	24.08	4.71	22.84	2.22	25.42	4.28	25.07	4.02

10

11 **Table 2:** Statistics in parameters measured at baseline and 12 months later.

Parameters	Mean G1	SD	Mean G2	SD	Mean G3	SD	Mean G4	SD
BMD L1-L4 (start)	0,812	0,08771	0,768	0,1032	0,87	0,1074	0,832	0,1556
BMD L1-L4 (end)	0,821	0,08737	0,8	0,104	0,856	0,1144	0,84	0,1209
BMD NECK LEFT (start)	0,64	0,07427	0,6	0,06343	0,655	0,08191	0,608	0,08176




Figure 1
162x104 mm (DPI)